Datasheet for ABIN1049145

anti-NR5A1 antibody (Internal Region)

2 Images

Go to Product page

Overview

Quantity:	50 µg
Target:	NR5A1
Binding Specificity:	Internal Region
Reactivity:	Human, Monkey, Bat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR5A1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Immunogen:	Synthetic 17 amino acid peptide from internal region of human NR5A1. Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Bat (100%), Elephant, Panda, Dog, Bovine, Rabbit, Horse, Pig (94%), Mouse, Rat, Hamster (88%), Turkey, Chicken, Platypus, Xenopus (82%). Type of Immunogen: Synthetic peptide
Specificity:	Human NR5A1. BLAST analysis of the peptide immunogen showed no homology with other human proteins, except TRIM23 (47 %).
Predicted Reactivity:	Percent identity with other species by BLAST analysis: Human, Gorilla, Gibbon, Monkey, Marmoset, Bat (100%) Elephant, Panda, Dog, Bovine, Rabbit, Horse, Pig (94%) Mouse, Rat,

Product Details

	Hamster (88%) Turkey, Chicken, Platypus, Xenopus (82%).
Purification:	Immunoaffinity purified

Target Details

Target:	NR5A1
Alternative Name:	NR5A1 / SF1 (NR5A1 Products)
Background:	Name/Gene ID: NR5A1 Subfamily: NR5 Fushi tarazu-like Family: NHR Synonyms: NR5A1, AD4BP, Adrenal 4 binding protein, Adrenal 4-binding protein, ELP, Steroidogenic factor-1, SRXY3, Steroidogenic factor 1, Nuclear receptor AdBP4, Nuclear receptor elp, POF7, SF-1, SPGF8, Steroid hormone receptor Ad4BP, STF-1, FTZ1, FTZF1, Fushi tarazu factor homolog 1
Gene ID:	2516
Pathways:	Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway, Maintenance of Protein Location

Application Details

Application Notes:	Approved: IHC, IHC-P (2 - 18 µg/mL)
Comment:	Target Species of Antibody: Human
Assay Procedure:	**The IHC-pro Immunohistochemistry Protocol** **Tissue Preparation** Formalin fixation and embedding in paraffin wax **Tissue Sectioning** Make 4-µm sections and place on pre-cleaned and charged microscope slides. Heat in a tissue-drying oven for 45 minutes at 60°C Deparaffinization Wash slides in 3 changes of xylene – 5 minutes each at room temperature.

Rehydration

Wash slides in 3 changes of 100% alcohol – 3 minutes each at room temperature.

Wash slides in 2 changes of 95% alcohol – 3 minutes each at room temperature.

Wash slides in 1 change of 80% alcohol – 3 minutes at room temperature.

Rinse slides in gentle running distilled water – 5 minutes at room temperature.

Antigen retrieval

Steam slides in 0.01 M sodium citrate buffer, pH 6.0 at 99-100°C - 20 minutes

Remove from heat and let stand at room temperature in buffer - 20 minutes

Rinse in 1X TBS with Tween (TBST) – 1 minute at room temperature.

Immunostaining

Do not allow tissues to dry at any time during the staining procedure.

Apply a universal protein block – 20 minutes at room temperature.

Drain protein block from slides, apply diluted primary antibody – 45 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply a biotinylated secondary antibody (specific to the host of the primary antibody) - 30 minutes at room temperature.

Rinse slides 1X TBST – 1 minute at room temperature.

Apply alkaline phosphatase streptavidin – 30 minutes at room temperature.

Rinse slides in 1X TBST - 1 minute at room temperature.

Apply alkaline phosphatase chromogen substrate – 30 minutes at room temperature.

Wash slides in distilled water – 1 minute at room temperature.

Dehydrate

This method should only be used if the chromogen substrate is alcohol insoluble.

Wash slides in 2 changes of 80% alcohol – 1 minute each at room temperature.

Wash slides in 2 changes of 95% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of 100% alcohol – 1 minute each at room temperature.

Wash slides in 3 changes of xylene – 1 minute each at room temperature.

Apply coverslip

Restrictions:	For Research Use only

Copyright antibodies-online. All rights reserved.

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, less than 0.1 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Aliquot and store undiluted at -20°C or below for up to 1 year. Can be stored undiluted at 4°C for up to 1 month. Avoid freeze-thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Anti-NR5A1 antibody ABIN1049145 IHC staining of human ovary. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.



Immunohistochemistry

Image 2. Anti-NR5A1 antibody IHC of human ovary. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval.