



Datasheet for ABIN106405
anti-RAD9A antibody (pSer1260)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	RAD9A
Binding Specificity:	AA 1249-1263, pSer1260
Reactivity:	Saccharomyces cerevisiae
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAD9A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to phosphorylated form of aa 1249-1263 of 1309 of yeast Rad9 protein conjugated to KLH.
Isotype:	IgG
Characteristics:	Concentration Definition: by UV absorbance at 280 nm

Target Details

Target:	RAD9A
Alternative Name:	Rad9 (RAD9A Products)
Background:	Rad9 is required for the MEC1/TEL1-dependent activation of Saccharomyces cerevisiae DNA damage checkpoint pathways mediated by Rad53 and Chk1. DNA damage induces Rad9

Target Details

phosphorylation, and Rad53 specifically associates with phosphorylated Rad9. Cells have evolved multiple strategies for tolerating genomic damage. The most important of these are numerous repair systems that remove or bypass potentially mutagenic DNA lesions. Another cellular strategy is to delay cell-cycle transitions at multiple points. The genetic control of these delays, termed 'checkpoints', was first established in budding yeast where it was shown that the RAD9 gene functions in G2/M arrest after irradiation with X-rays. Subsequently, it has become clear that Rad9 also functions at the G1/S, intra-S and mid-anaphase checkpoints. Defects in checkpoint regulation can lead to genome instability and, in higher eukaryotes, neoplastic transformation. Rad9 also controls the transcriptional induction of a DNA damage regulon (DDR). Rad9 may also have a pro-apoptotic function. This is suggested in that Rad9 from *Schizosaccharomyces pombe* (SpRad9) contains a group of amino acids with similarity to the Bcl-2 homology 3 death domain, which is required for SpRad9 interaction with human Bcl-2 and apoptosis induction in human cells. Overexpression of Bcl-2 in *S. pombe* inhibits cell growth independently of rad9, but enhances resistance of rad9-null cells to methyl methanesulfonate, ultraviolet and ionizing radiation. Rad9 conveys the checkpoint signal by activating Rad53p and Chk1p; is hyperphosphorylated by Mec1p and Tel1p; and is a potential Cdc28p substrate. Mature yeast Rad9 is reported to have an apparent molecular weight of ~148kDa. The human homolog is reported at 48.5 kDa.

Synonyms: Cell cycle checkpoint control protein antibody, Cell cycle checkpoint control protein RAD9A antibody, DNA repair exonuclease rad9 homolog A antibody

Gene ID: 851803, 4759022

UniProt: [Q99638](#)

Pathways: [Positive Regulation of Response to DNA Damage Stimulus](#)

Application Details

Application Notes: This phospho specific polyclonal antibody was tested by immunoblotting and ELISA. Data from both immunoblotting and ELISA indicate the antibody is reactive with the phosphorylated form of the immunizing peptide and minimally reactive with the non-phosphorylated form of the immunizing peptide. Immunoblotting detects yeast Rad9 protein. No reactivity is expected against the human or mouse analogs of RAD9. Reactivity against RAD9 from other sources is unknown. Cross reactivity may occur with auto-phosphorylated Rad53 kinase. Although not tested, this antibody is likely functional by IHC and IP. This product has been assayed against 0.1 µg of phosphorylated peptide (pS1260) in a standard capture ELISA using TMB (3,3',5,5'-Tetramethylbenzidine) as a substrate for 30 minutes at room temperature. A working dilution

Application Details

of 1:5,000 is suggested for this product. Minimal reactivity was detected against the non-phosphorylated form (S1260) of the immunizing peptide. This antibody appears to be specific for the active form (phosphorylated) of the protein. Dilute the antibody 1:100 to 1:500 for immunoblotting. Researchers should determine optimal titers for other applications.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.41 mg/mL

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Images

Schematic summary of the DNA replication and DNA damage checkpoints in *S. cerevisiae*.





Image 1. Checkpoints are mechanisms that impose delays in the cell cycle in response to DNA damage or defects in DNA replication, to ensure that mitotic transmission is error-free. Failure to delay the cell cycle in the presence of damage converts an easily repairable DNA lesion into one far more deleterious, provoking genomic instability or cell death. This figure shows a summary of our current knowledge about the DNA damage checkpoints in yeast. Genetic analysis of the pathway has allowed classification of its components into “Sensors”, which detect different sorts of damage, “Signal transducers” which are signal-integrating kinases, and “Targets” which carry out the essential functions of suppressing progress through the cell cycle (i.e. inducing repair genes and preventing late origin firing or sister chromatid segregation. Contributed by C. Frei



and K. Shimada, laboratory of S. Gasser, U. Geneva.

Western Blotting

Image 2. Affinity purified phospho-specific antibody to yeast Rad9 at pS1260 was used at a 1:200 dilution incubated overnight at 4° C to detect Rad9 by Western blot. Lanes were loaded with 50 ng each of recombinant GST fusion protein containing *S. cerevisiae* Rad9 (aa 991-1309 ~60 kDa) on a 4-20% Criterion gel for SDS-PAGE as follows: Lane 1 - non-phosphorylated wild type yeast Rad9, Lane 2 - in vitro phosphorylated wild type yeast Rad9, Lane 3 - non-phosphorylated S1129A/S1260A double mutant Rad9, Lane 4 - in vitro phosphorylated S1129A/S1260A double mutant. Phosphorylation of Rad9 was by treatment with ATP and Rad53 kinase. Rad53 kinase autophosphorylates and appears cross reactive as it is detected on the blot as 90 and 110 kDa bands (asterisk). Detection occurred using a 1:5,000 dilution of 800 conjugated Donkey anti-Rabbit IgG (code # 611-732-127) for 1h at room temperature. LICOR's Infrared Imaging System was used to scan and process the image. Other detection systems will yield similar results.