



Datasheet for ABIN1881328  
**anti-EXOSC2 antibody (N-Term)**



[Go to Product page](#)

1 Image

2 Publications

### Overview

Quantity:	400 µL
Target:	EXOSC2
Binding Specificity:	AA 1-30, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOSC2 antibody is un-conjugated
Application:	Western Blotting (WB)

### Product Details

Immunogen:	This EXOSC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human EXOSC2.
Isotype:	Ig Fraction
Purification:	This antibody is purified through a protein A column, followed by peptide affinity purification.

### Target Details

Target:	EXOSC2
Alternative Name:	EXOSC2 ( <a href="#">EXOSC2 Products</a> )
Background:	Non-catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the nucleus, the RNA exosome complex is involved in proper maturation of stable RNA species

## Target Details

---

such as rRNA, snRNA and snoRNA, in the elimination of RNA processing by-products and non-coding 'pervasive' transcripts, such as antisense RNA species and promoter-upstream transcripts (PROMPTs), and of mRNAs with processing defects, thereby limiting or excluding their export to the cytoplasm. The RNA exosome may be involved in Ig class switch recombination (CSR) and/or Ig variable region somatic hypermutation (SHM) by targeting AICDA deamination activity to transcribed dsDNA substrates. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. The catalytic inactive RNA exosome core complex of 9 subunits (Exo-9) is proposed to play a pivotal role in the binding and presentation of RNA for ribonucleolysis, and to serve as a scaffold for the association with catalytic subunits and accessory proteins or complexes. EXOSC2 as peripheral part of the Exo-9 complex stabilizes the hexameric ring of RNase PH-domain subunits through contacts with EXOSC4 and EXOSC7.

Molecular Weight:	32789
NCBI Accession:	<a href="#">NP_055100</a>
UniProt:	<a href="#">Q13868</a>
Pathways:	<a href="#">SARS-CoV-2 Protein Interactome</a>

## Application Details

---

Application Notes:	WB: 1:1000
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Buffer:	Purified polyclonal antibody supplied in PBS with 0.09 % (W/V) sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Expiry Date:	6 months

## Publications

Product cited in:

Dai, Liu, Liu, Zhang, Wang, Jin, Qian, Wang, Zhao, Wu, Xiong, Chang, Sun, Yang, Hoffman, Liu: "Anti-metastatic Efficacy of Traditional Chinese Medicine (TCM) Ginsenoside Conjugated to a VEGFR-3 Antibody on Human Gastric Cancer in an Orthotopic Mouse Model." in: **Anticancer research**, Vol. 37, Issue 3, pp. 979-986, (2017) ([PubMed](#)).


Irrthum, Karkkainen, Devriendt, Alitalo, Vikkula: "Congenital hereditary lymphedema caused by a mutation that inactivates VEGFR3 tyrosine kinase." in: **American journal of human genetics**, Vol. 67, Issue 2, pp. 295-301, (2000) ([PubMed](#)).

Galland, Karamysheva, Pebusque, Borg, Rottapel, Dubreuil, Rosnet, Birnbaum: "The FLT4 gene encodes a transmembrane tyrosine kinase related to the vascular endothelial growth factor receptor." in: **Oncogene**, Vol. 8, Issue 5, pp. 1233-40, (1993) ([PubMed](#)).

Pajusola, Aprelikova, Korhonen, Kaipainen, Pertovaara, Alitalo, Alitalo: "FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines." in: **Cancer research**, Vol. 52, Issue 20, pp. 5738-43, (1992) ([PubMed](#)).

Galland, Karamysheva, Mattei, Rosnet, Marchetto, Birnbaum: "Chromosomal localization of FLT4, a novel receptor-type tyrosine kinase gene." in: **Genomics**, Vol. 13, Issue 2, pp. 475-8, (1992) ([PubMed](#)).

## Images



### Western Blotting

**Image 1.** EXOSC2 Antibody (N-term) (ABIN1881328 and ABIN2843634) western blot analysis in Jurkat cell line lysates (35 µg/lane). This demonstrates the EXOSC2 antibody detected the EXOSC2 protein (arrow).