



[Go to Product page](#)

Datasheet for ABIN2175303
anti-HCST antibody (AA 19-48) (PE)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | HCST |
| Binding Specificity: | AA 19-48 |
| Reactivity: | Human, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This HCST antibody is conjugated to PE |
| Application: | Western Blotting (WB), Flow Cytometry (FACS) |

Product Details

| | |
|-------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human HCST |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Rat |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | HCST |
| Alternative Name: | HCST (HCST Products) |
| Background: | Synonyms: DAP10, KAP10, PIK3AP, Hematopoietic cell signal transducer, DNAX-activation protein 10, Membrane protein DAP10, Transmembrane adapter protein KAP10, HCST, |

Target Details

UNQ587/PRO1157

Background: Transmembrane adapter protein which associates with KLRK1 to form an activation receptor KLRK1-HCST in lymphoid and myeloid cells, this receptor plays a major role in triggering cytotoxicity against target cells expressing cell surface ligands such as MHC class I chain-related MICA and MICB, and UL16-binding proteins (ULBPs), these ligands are up-regulated by stress conditions and pathological state such as viral infection and tumor transformation. Functions as docking site for PI3-kinase PIK3R1 and GRB2. Interaction of ULBPs with KLRK1-HCST triggers calcium mobilization and activation of the PIK3R1, MAP2K/ERK, and JAK2/STAT5 signaling pathways. Both PIK3R1 and GRB2 are required for full KLRK1-HCST-mediated activation and ultimate killing of target cells. In NK cells, KLRK1-HCST signaling directly induces cytotoxicity and enhances cytokine production initiated via DAP12/TYROBP-associated receptors. In T-cells, it provides primarily costimulation for TCR-induced signals. KLRK1-HCST receptor plays a role in immune surveillance against tumors and is required for cytolysis of tumors cells, indeed, melanoma cells that do not express KLRK1 ligands escape from immune surveillance mediated by NK cells.

Gene ID: 10870

UniProt: [Q9UBK5](#)

Application Details

Application Notes: FCM 1:20-100

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months