



## Datasheet for ABIN2782428 anti-NCF4 antibody (N-Term)

1 Image



[Go to Product page](#)

### Overview

Quantity:	100 µL
Target:	NCF4
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Dog, Rabbit, Horse, Guinea Pig, Pig, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NCF4 antibody is un-conjugated
Application:	Western Blotting (WB)

### Product Details

Immunogen:	The immunogen is a synthetic peptide directed towards the N terminal region of human NCF4
Sequence:	SKYLIYRRYR QFHALQSKLE ERFGPDSKSS ALACTLPTLP AKVYVGVKQE
Predicted Reactivity:	Cow: 85%, Dog: 86%, Guinea Pig: 79%, Horse: 86%, Human: 100%, Mouse: 92%, Pig: 92%, Rabbit: 85%, Rat: 92%
Characteristics:	This is a rabbit polyclonal antibody against NCF4. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified

### Target Details

Target:	NCF4
---------	------

## Target Details

---

Alternative Name:	NCF4 ( <a href="#">NCF4 Products</a> )
Background:	<p>NCF4 is a cytosolic regulatory component of the superoxide-producing phagocyte NADPH-oxidase, a multicomponent enzyme system important for host defense. It interacts primarily with neutrophil cytosolic factor 2 (NCF2/p67-phox) to form a complex with neutrophil cytosolic factor 1 (NCF1/p47-phox), which further interacts with the small G protein RAC1 and translocates to the membrane upon cell stimulation. This complex then activates flavocytochrome b, the membrane-integrated catalytic core of the enzyme system. The PX domain of this protein can bind phospholipid products of the PI(3) kinase, which suggests its role in PI(3) kinase-mediated signaling events. The phosphorylation of this protein was found to negatively regulate the enzyme activity. Alternatively spliced transcript variants encoding distinct isoforms have been observed. The protein encoded by this gene is a cytosolic regulatory component of the superoxide-producing phagocyte NADPH-oxidase, a multicomponent enzyme system important for host defense. This protein is preferentially expressed in cells of myeloid lineage. It interacts primarily with neutrophil cytosolic factor 2 (NCF2/p67-phox) to form a complex with neutrophil cytosolic factor 1 (NCF1/p47-phox), which further interacts with the small G protein RAC1 and translocates to the membrane upon cell stimulation. This complex then activates flavocytochrome b, the membrane-integrated catalytic core of the enzyme system. The PX domain of this protein can bind phospholipid products of the PI(3) kinase, which suggests its role in PI(3) kinase-mediated signaling events. The phosphorylation of this protein was found to negatively regulate the enzyme activity. Alternatively spliced transcript variants encoding distinct isoforms have been observed.</p>
	Alias Symbols: MGC3810, NCF, P40PHOX, SH3PXD4
	Protein Interaction Partner: NCF2, UBC, CORO1A, MSN, CYBB, NCF1, CYBA, TXN, PRKCD, XRCC6, PRKDC,
	Protein Size: 348
Molecular Weight:	39 kDa
Gene ID:	4689
NCBI Accession:	<a href="#">NM_013416</a> , <a href="#">NP_038202</a>
UniProt:	<a href="#">A8K4F9</a>

## Application Details

---

Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 348 AA

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

## Images

---

