



[Go to Product page](#)

Datasheet for ABIN3091919
CSK Protein (AA 2-450) (His tag)

Overview

Quantity:	1 mg
Target:	CSK
Protein Characteristics:	AA 2-450
Origin:	Human
Source:	Insect Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CSK protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Crystallization (Crys)

Product Details

Sequence: SAIQAAWPSG TECIAKYNFH GTAEQDLPFC KGDVLTIVAV TKDPNWKAK NKVREGIIP
ANYVQKREGV KAGTKLSLMP WFHGKITREQ AERLLYPET GLFLVRESTN YPGDYTL CVS
CDGKVEHYRI MYHASKLSID EEVYFENLMQ LVEHYTSDAD GLCTRLIKPK VMEGTVAAQD
EFYRSGWALN MKELKLLQTI GKGEFGDVML GDYRGNKVAV KCIKNATAQ AFLAEASVMT
QLRHSNLVQL LGVIVEEKGK LYIVTEYMAK GSLVDYLRSR GRSVLGGDCL LKFSLDVCEA
MEYLEGNNFV HRDLAARNVL VSEDNVAKVS DFGLTKEASS TQDTGKLPVK WTAPEALREK
KFSTKSDVWS FGILLWEIYS FGRVPYPRIP LKDVVPRVEK GYKMDAPDGC PPAVYEV MKN
CWHLDAAMRP SFLQLREQL HIKTHELHL

Sequence without tag. Tag location is at the discretion of the manufacturer. If you have a special request, please contact us.

- Characteristics:
- Made in Germany - from design to production - by highly experienced protein experts.
 - Human CSK Protein (raised in Insect Cells) purified by multi-step, protein-specific process to

Product Details

ensure crystallization grade.

- State-of-the-art algorithm used for plasmid design (Gene synthesis).

This protein is a made to order protein and will be made for the first time for your order. Our experts in the lab will ensure that you receive a correctly folded protein.

The big advantage of ordering our made-to-order proteins in comparison to ordering custom made proteins from other companies is that there is no financial obligation in case the protein cannot be expressed or purified.

In the unlikely event that the protein cannot be expressed or purified we do not charge anything (other companies might charge you for any performed steps in the expression process for custom-made proteins, e.g. fees might apply for the expression plasmid, the first expression experiments or purification optimization).

When you order this made-to-order protein you will only pay upon receipt of the correctly folded protein. With no financial risk on your end you can rest assured that our experienced protein experts will do everything to make sure that you receive the protein you ordered.

The concentration of our recombinant proteins is measured using the absorbance at 280nm. The protein's absorbance will be measured in several dilutions and is measured against its specific reference buffer.

The concentration of the protein is calculated using its specific absorption coefficient. We use the Expasy's protparam tool to determine the absorption coefficient of each protein.

Purification:	Two step purification of proteins expressed in baculovirus infected SF9 insect cells: <ol style="list-style-type: none">1. In a first purification step, the protein is purified from the cleared cell lysate using three different His-tag capture materials: high yield, EDTA resistant, or DTT resistant. Eluate fractions are analyzed by SDS-PAGE.2. Protein containing fractions of the best purification are subjected to second purification step through size exclusion chromatography. Eluate fractions are analyzed by SDS-PAGE and Western blot.
---------------	---

Purity:	>95 % as determined by SDS PAGE, Size Exclusion Chromatography and Western Blot.
---------	--

Sterility:	0.22 µm filtered
------------	------------------

Endotoxin Level:	Protein is endotoxin free.
------------------	----------------------------

Grade:	Crystallography grade
--------	-----------------------

Target Details

Target:	CSK
---------	-----

Alternative Name:	CSK (CSK Products)
-------------------	--------------------------------------

Target Details

Background:	Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T-cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK. {ECO:0000269 PubMed:1639064, ECO:0000269 PubMed:9281320}.
Molecular Weight:	51.5 kDa Including tag.
UniProt:	P41240
Pathways:	TCR Signaling , EGFR Signaling Pathway , Cell-Cell Junction Organization , CXCR4-mediated Signaling Events

Application Details

Application Notes:	In addition to the applications listed above we expect the protein to work for functional studies as well. As the protein has not been tested for functional studies yet we cannot offer a guarantee though.
Comment:	In cases in which it is highly likely that the recombinant protein with the default tag will be insoluble our protein lab may suggest a higher molecular weight tag (e.g. GST-tag) instead to increase solubility. We will discuss all possible options with you in detail to assure that you receive your protein of interest.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	100 mM NaCL, 20 mM Hepes, 10% glycerol. pH value is at the discretion of the manufacturer.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-80 °C
Storage Comment:	Store at -80°C.

Handling

Expiry Date: Unlimited (if stored properly)